

us—and other readers—whether gender independently influenced clinical outcome.

**\*Tina S. Poulsen, MD**  
**Anders Kastrup, MD**  
**Hans Mickley, MD, PhD**

\*Department of Cardiology  
 Odense University Hospital  
 Sdr. Boulevard 29  
 DK-5000 Odense C  
 Denmark  
 E-mail: [tina.poulsen@ouh.fyns-amt.dk](mailto:tina.poulsen@ouh.fyns-amt.dk)

doi:10.1016/j.jacc.2004.11.019

## REFERENCES

1. Chen WH, Lee PY, Ng W, et al. Aspirin resistance is associated with a high incidence of myonecrosis after non-urgent percutaneous coronary intervention despite clopidogrel pretreatment. *J Am Coll Cardiol* 2004;43:1122–6.
2. King KM, Ghali WA, Faris PD, et al. Sex differences in outcomes after cardiac catheterization. Effect modification by treatment strategy and time. *JAMA* 2004;291:1220–5.
3. Vaccarino V, Abramson JL, Veledar E, et al. Sex-based differences in early mortality after myocardial infarction. *N Engl J Med* 1999;341:217–25.

## REPLY

We would like to thank Dr. Poulsen and colleagues for their interest in our study (1). The prevalence of female gender with or without creatine kinase-myocardial band (CK-MB) elevation after percutaneous coronary intervention (PCI) was not different between the two groups (31.1% vs. 21.7%;  $p = 0.22$ ). Gender was entered in the multivariate analysis, and it turned out not to be an independent predictor of CK-MB elevation.

Women undergoing balloon angioplasty had a higher incidence of early adverse cardiac events in both early (2) and contemporary series (3). In the current era of predominant stent implantation for PCI, mixed results have been demonstrated (4–6). However, our data concur with recent studies specifically addressing post-PCI myonecrosis, in which female gender is not demonstrated to be an independent correlate of post-PCI myonecrosis (7–9).

**\*Wai-Hong Chen, MBBS**  
**Pui-Yin Lee, MBBS**  
**William Ng, MBBS**  
**Hung-Fat Tse, MD**  
**Chu-Pak Lau, MD**

\*Division of Cardiology  
 Department of Medicine  
 The University of Hong Kong  
 Queen Mary Hospital  
 Hong Kong  
 China  
 E-mail: [whchen@hku.hk](mailto:whchen@hku.hk)

doi:10.1016/j.jacc.2004.11.018

## REFERENCES

1. Chen WH, Lee PY, Ng W, et al. Aspirin resistance is associated with a high incidence of myonecrosis after non-urgent percutaneous coronary intervention despite clopidogrel pretreatment. *J Am Coll Cardiol* 2004;43:1122–6.
2. Cowley MJ, Mullin SM, Kelsey SF, et al. Sex differences in early and long-term results of coronary angioplasty in the NHLBI PTCA registry. *Circulation* 1985;71:90–7.
3. Kelsey SF, James M, Hloubkov R, et al. Results of percutaneous transluminal coronary angioplasty in women. 1985–1986 National Heart, Lung, and Blood Institute's Coronary Angioplasty Registry. *Circulation* 1993;87:720–7.
4. Mehilli J, Kastrati A, Dirschinger J, et al. Differences in prognostic factors and outcomes between women and men undergoing coronary artery stenting. *JAMA* 2000;284:1799–805.
5. Watanabe CT, Maynard C, Ritchie JL. Comparison of short-term outcomes following coronary artery stenting in men versus women. *Am J Cardiol* 2001;88:848–52.
6. Trabattini D, Bartorelli AL, Montorsi P, et al. Comparison of outcomes in women and men treated with coronary stent implantation. *Catheter Cardiovasc Interv* 2003;58:20–8.
7. Kini A, Marmur JD, Kini S, et al. Creatine kinase-MB elevation after coronary intervention correlates with diffuse atherosclerosis, and low-to-medium level elevation has a benign clinical course. *J Am Coll Cardiol* 1999;34:663–71.
8. Stone GW, Mehran R, Gangas G, et al. Differential impact on survival of electrocardiographic Q-wave versus enzymatic myocardial infarction after percutaneous intervention: a device-specific analysis of 7,147 patients. *Circulation* 2001;104:642–7.
9. Kini AS, Lee P, Mitre CA, et al. Postprocedure chest pain after coronary stenting: implications on clinical restenosis. *J Am Coll Cardiol* 2003;41:33–8.